

LA-UR-21-29859

Approved for public release; distribution is unlimited.

Title: Dual n-gamma Measurement of the $^{235}\text{U}(n,3n)$ Cross Section

Author(s): Kelly, Keegan John
Mosby, Michelle A.
Kawano, Toshihiko

Intended for: 2021 LDRD-DR Pitch Town Hall

Issued: 2021-10-05

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Dual n - γ Measurement of the $^{235}\text{U}(n,3n)$ Cross Section

Keegan Kelly (P-3, Presenting),
Michelle Mosby (XTD-NTA), Toshihiko Kawano (T-2)

October 5, 2021

Background

Context/Timeliness

- See Impact/Strategic Relevance section.
- The detector array intended for use in this proposal will reach maturity at exactly the time of the proposed ^{235}U measurement.

Knowledge Gap

- $^{235}\text{U}(n,3n)$ cross section is poorly known.
- Data capable of yielding the proposed types of cross sections will be available, but the methods of extracting need to be developed.

Innovation

- Applying dual n - γ detection analysis with newly-available CLYC detectors.
- Array of CLYC detectors is currently being developed and built under LDRD project 20210329ER, which will reach the full- or near-full-scale design during the second year of the proposed DR project. This is when the ^{235}U measurement would occur.
- Fully map the n - γ correlation space from $(n,2n)$ and $(n,3n)$ reactions, which has never been measured before.
- New, independent method of extracting highly-desirable cross sections.

Research Approach

Hypothesis/Science Question

- Can we develop methods to extract $(n,2n)$ and $(n,3n)$ cross sections from dual n - γ data obtained with a newly-developed highly-segmented array of CLYC detectors?

Strategic Collaborators

- Keegan Kelly, PI (P-3)
- Michelle Mosby, Co-PI (XTD-NTA)
- Toshihiko Kawano, Co-PI (T-2)

Methods

- Year 1: Leverage ^{197}Au and ^{238}U dual n - γ data collected with a mostly-complete CLYC detector array from LDRD project 20210329ER to develop methods to extract $(n,2n)$ and $(n,3n)$ cross sections, and define required experimental parameters for the ^{235}U measurement
- Year 2: Perform the ^{235}U measurement with the complete or near-complete CLYC detector array from project 20210329ER, taking experimental considerations from yr. 1 into account, and extract the $^{235}\text{U}(n,3n)$ cross section.
- Year 3: Perform new evaluation of $^{235}\text{U}+n$ cross sections, and assess impact.

Strategic Impact



Pillar

- Weapons Systems

Transition Plan

- Following the conclusion of this LDRD project, we will have developed a capability which should be attractive to OES.

Impact/Strategic Relevance to LANL

- The capability to do these types of measurements is important to the weapons community. Details are available on request.
- LA-CP-21-00199, LA-CP-21-00273, LA-CP-21-00292, LA-CP-21-00413, and more.